Dilution



CRISPR-Cas9 SP Rabbit mAb [hoiT]

Cat NO. :A26938

Information:

Applications detail:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC ICC/IF FC	Streptococcus	Q99ZW2	158kDa	Rabbit	IgG	50ul,100ul,200ul
	pyogenes					

Application

	WB	1:1000-2000
	IHC	1:100
	ICC/IF	1:100
	The optimal dilutions should be	determined by the end user
Conjugate:		
UnConjugate		
Form:		
Liquid		
sensitivity:		
Endogenous		
Purification:		
Affinity-chromatography		
Specificity:		
Antibody is produced by immunizing anim	nals with A synthesized peptide de	erived from human CRISPR-Cas9 SF
Storage buffer and conditions	5 :	
Antibody store in 10 mM PBS, 0.5mg/ml B	SSA, 50% glycerol (buffer) .	
Shipped at 4°C. Store at-20°C or -80°C.		
Products are valid for one natural year of	receipt. Avoid repeated freeze / tl	haw cycles.
Tissue specificity:		
Subcellular location:		
Cascolidiai location.		
Function:		

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

CRISPR-Cas9 SP Rabbit mAb [hoiT] Images



View more information on http://naturebios.com



IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.