

GLUT2 Rabbit mAb [07NX]

Cat NO. :A13235

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	P11168	57 kda	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human GLUT2.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Liver, insulin-producing beta cell, small intestine and kidney...

Subcellular location:

Cell membrane, Multi-pass membrane protein.

Function:

Facilitative hexose transporter that mediates the transport of glucose and fructose (PubMed:8027028, PubMed:16186102, PubMed:23396969, PubMed:28083649). Likely mediates the bidirectional transfer of glucose across the plasma membrane of hepatocytes and is responsible for uptake of glucose by the beta cells,may comprise part of the glucose-sensing mechanism of the beta cell (PubMed:8027028). May also participate with the Na(+)/glucose cotransporter in the transcellular transport of glucose in the small intestine and kidney (PubMed:3399500). Also able to mediate the transport of dehydroascorbate (PubMed:23396969)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

GLUT2 Rabbit mAb [07NX] Images



Western blot (SDS PAGE) analysis of extracts from Human fetal liver tissue lyaste.using GLUT2 Rabbit mAb [07NX] at dilution of 1:1000 incubated at $4\,^{\circ}\mathrm{C}$ over night

View more information on http://naturebios.com