

CDKN2A/p16INK4a Rabbit mAb [7VX4]

Cat NO. :A47185

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	н	P42771	17	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

ICC/IF

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

 $Antibody \ is \ produced \ by \ immunizing \ animals \ with \ a \ synthetic \ peptide \ at \ the \ sequence \ of \ human \ CDKN2A/p16INK4a.$

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Widely expressed but not detected in brain or skeletal muscle. Isoform 3 is pancreas-specific..

Subcellular location:

Cytoplasm. Nucleus.

Function:

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6.

This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

CDKN2A/p16INK4a Rabbit mAb [7VX4] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells.Using CDKN2A/p16INK4a Rabbit mAb [7VX4] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com