

DNA polymerase mu Rabbit mAb [3RLS]

Cat NO. :A66214

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	Q9NP87	55 kDa	Rabbit	IgG	100ul,200ul

Applications detail:	Application	Dilution		
	wв	1:1000-2000		
	The optimal dilutions should be determined by the end user			

Conjugate:	
UnConjugate	
Form:	
Liquid	
sensitivity:	
Endogenous	
Purification:	
Protein A purification	

Specificity:
Antibody is produced by immun

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human DNA polymerase mu

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed in a number of tissues. Abundant in thymus.

Subcellular location:

Nucleus.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Gap-filling polymerase involved in repair of DNA double-strand breaks by non-homologous end joining (NHEJ). Participates in immunoglobulin (Ig) light chain gene rearrangement in V(D)J recombination..

Validation Data:

DNA polymerase mu Rabbit mAb [3RLS] Images



Western blot (SDS PAGE) analysis of extracts from 293T cells. Using DNA polymerase muRabbit mAb [3RLS] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com