

MGP Mouse mAb[C8G3]

Cat NO. :A81253

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size	
WB,IHC	H,M	P08493	12kDa	Mouse	IgG	100ul,200ul	

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

					te:		
	\sim 10		~	~	-	-	
\mathbf{u}	UH	IIU	u	а	LE	=	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human MGP.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Secreted.

Function:

Associates with the organic matrix of bone and cartilage. Thought to act as an inhibitor of bone formation.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

MGP Mouse mAb[C8G3] Images



Western blot (SDS PAGE) analysis of extracts from human placenta tissue. Using MGP Mouse mAb IgG [C8G3] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com